

Running Head: Affect, Dual Processing & Health

Affect, Dual-processing, Developmental Psychopathology & Health Behaviors

Reinout W. Wiers¹
Kristen G. Anderson²
Bram Van Bockstaele¹
Elske Salemink¹
Bernhard Hommel³

1. Addiction, Development and Psychopathology (ADAPT)-lab, Department of Psychology, University of Amsterdam & Research Priority Area Yield
2. Adolescent Health Research Program, Department of Psychology, Reed College
3. Leiden University.

Chapter to appear in “Affective Determinants of Health Behavior” Editors: David Williams, Ryan Rhodes, and Mark Conner. NY: Oxford University Press.

Final In Press Version 

Abstract (120-150 words)

In this chapter we discuss dual-process models of (health) behaviors, both regarding their recent criticisms and regarding implications for health interventions. We agree with critics that impulsive and reflective processes should not be equated with specific brain processes, but that psychological processes are emergent properties of the dynamic, in time unfolding interplay between different neural systems. We maintain that at a psychological level of description, these models can still be useful to understand challenges to health behaviors and possible interventions. Affective processes can influence impulsive decision making in health, but also reflective processes, when they concern affectively relevant goals. Cognitive training methods, including Cognitive Bias Modification and training of Executive Control, have shown some success in changing health-behaviors, but a critical variable for long-term success appears to be motivation to change.

Keywords: Dual Process Models, Affect, Health, Development, Cognitive Bias Modification

Introduction

In this chapter, we develop a broad theoretical framework to account for healthy and unhealthy behaviors and related psychopathology from a developmental perspective. We start with a discussion of dual process models, and will argue that from a neurocognitive perspective, dual (or multiple) processes are an emergent property of unfolding and interacting brain processes. Although dual process models have been criticized, we argue that these models can still be useful at the psychological level of description. We then turn to broad developmental and personality inspired models including differences in temperament and socialization in relation to the development of individual differences in outcomes of health-relevant behaviors. The general heuristic model (Gladwin, Figner, Crone, & Wiers, 2011), is illustrated with a description of experiments, on the interplay between bottom-up and top-down processes in relation to health behaviors. In the second part of the chapter, we discuss interventions from this perspective.

Affect and Cognition, Dual-Process Models and Beyond

Dual process models have been *en vogue* the past decades in psychology (Evans, 2008; Smith & DeCoster, 2000; Strack & Deutsch, 2004). They have ancient roots, emphasizing the struggle between reason and passion, or the heart and the head (Hofmann, Friese, & Strack, 2009). Many different names have been given to the different processing modes (sometimes called systems), we here use the terms “impulsive” and “reflective” processes (Strack & Deutsch, 2004). Note that this does not equate to unconscious and conscious, some problem-solving occurs unconsciously (Kounios & Beeman, 2014), and some seemingly rash impulsive behaviors may be entirely goal-directed (see for intriguing examples, Kopetz & Orehek, 2015). As the historical background suggests, affective influences have primarily been associated with impulsive processes and non-

affective influences with reflective processes. However, as emotions get a stronger cognitive component (e.g. shame, guilt), they may also fuel reflective processes to inhibit an impulse to prevent (anticipated) shame, regret, or guilt. Some theorists have emphasized different evolutionary origins, with impulsive processes representing evolutionary old processes and reflective processes evolutionary later additions (Evans, 2008; K. B. MacDonald, 2008); some have emphasized different developmental origins, with impulsive processes emerging earlier in life than reflective (effortful control) processes (Nigg, 2006; Rothbart & Posner, 2011); some have distinguished between associative (impulsive) processes and symbolic (reflective) processes (Gawronski & Bodenhausen, 2006; Strack & Deutsch, 2004); and some have distinguished between model-free (impulsive) vs. model-based (reflective) learning mechanisms (Ashby, Alfonso-Reese, Turken, & Waldron, 1998; Daw, Niv, & Dayan, 2005). In addition to general dual-process models, more specific models have been developed to account for the development of common mental disorders, such as addictive behaviors (Bechara, 2005; Wiers & Stacy, 2006), anxiety (Ouimet, Gawronski, & Dozois, 2009), and depression (Beevers, 2005). Dual-process models have also been adapted to account for health behaviors (Hofmann, Friese, & Wiers, 2008; Sheeran, Gollwitzer, & Bargh, 2013). Before we return to dual process models and health, it is important to consider some of the recent criticisms on dual process models, and to consider what can be maintained regarding a general model on cognitive-motivational processes involved in health psychology and related psychopathology.

Neurocognitive Processes and Dual Process Models

In an influential criticism of dual process/systems models, it was argued that there is no evidence for separable neural systems for impulsive and reflective processes (Keren & Schul, 2009). We agree to some extent with the criticisms on those dual

process models in which authors have equated specific brain systems with impulsive (or “reflexive”) processes (including “emotional” structures like the amygdala and striatum), and other areas, including parts of the prefrontal and parietal cortex as dealing exclusively with reflective processing (Heatherton & Wagner, 2011; Satpute & Lieberman, 2006), or in more broad, unspecified terms that different brain areas are correlated with impulsive and reflective processes (Evans & Stanovich, 2013). One problem is that this does not prove anything, many other models would be compatible with that observation (Keren, 2013).

But is this a fatal blow for dual process models? We argued it is not (Gladwin et al., 2011; Wiers, Gladwin, Hofmann, Salemink, & Ridderinkhof, 2013). What the criticism rules out are dual process models that propose a one-to-one relationship between mental processes and the underlying neurocognitive machinery, but alternatives are conceivable. For instance, the MIMIC (Multiple Indicators, Multiple Causes) model is based on the supervenience thesis (Kievit et al., 2011), according to which psychological constructs can be realized by underlying neural structures or processes in multiple ways. The neural indicators determine/constrain the psychological processes, without necessarily having a one-to-one relationship with them. This is in line with findings in cognitive neuroscience that the same neural structures are implicated in many psychological processes (see Kievit et al., 2011).

According to another perspective on the relationship between brain- and mental processes, the processes described in dual process models are emergent properties of the dynamic, in time unfolding interplay between different neural systems (Cunningham & Zelazo, 2007; Gladwin et al., 2011; Hazy, Frank, & O’Reilly, 2007). This perspective suggests that behavior that can be described according to a dual process model might actually arise from interactions among several neural systems, each with their own

computational objectives, and that even any one of the ‘processes’ reflects a dynamic interaction among these same systems (Frank, Cohen, & Sanfey, 2009). Importantly, recent computational models have demonstrated how symbolic processing, characteristic of reflective processing, can arise via dynamic interactions among multiple “loops” connecting the prefrontal cortex and basal ganglia (Collins & Frank, 2013; Kriete, Noelle, Cohen, & O’Reilly, 2013). According to these models, there can still be a first crude pass of information, which in extreme cases may already activate action-tendencies (“impulsive action”). However, typically more loops of information processing take place, which can be described as increasingly reflective *states* of processing (Gladwin et al., 2011). Clearly, this does not allow for specific neural structures (e.g. the dorsolateral prefrontal cortex) to only be involved in reflective processes and other structures (e.g., the amygdala) in impulsive processes (see for a similar argument regarding emotion and cognition, (Pessoa, 2008)).

Where does this leave us at the psychological level? At that level of description (i.e., models describing the interactions between psychological constructs, without equating them to underlying neural processes or systems), we would argue that dual process models could still be useful in generating testable predictions for understanding health-related processes. For example, in the case of health psychology, one paradoxical prediction of dual process models is that health prevention messages can backfire, especially when they are negatively phrased (e.g., “alcohol does not make you sexy”). The reason is that negation is a typical example of symbolic processing, and that the presentation with the two concepts (alcohol – sexy) can actually be strengthened by exposure, when over time the “not” part gets lost (Deutsch, Gawronski, & Strack, 2006; Krank, Ames, Grenard, Schoenfeld, & Stacy, 2010).

Development, Cognitive Processes, Psychopathology and Health

While numerous terms have been used in the literature to describe self-regulatory capacity, these functions generally represent our ability to exert control, inhibit, or regulate attention, behavior, and cognition (Ahadi & Rothbart, 1994; Eisenberg et al., 2013; Nigg, 2000), and are commonly contrasted with dispositional approach tendencies, impulsivity, or disinhibition (Gray, 1994; Whiteside & Lynam, 2003), with both changing substantively across development (Anderson & Briggs, 2016). Before age 2, attentional control is the predominant means of coping with emotional distress with signs of more classical behavioral regulation developing by age 3 (Anderson & Briggs, 2016; Eisenberg, Smith, & Spinrad, 2010). Self-control capacity increases dramatically in the preschool years and continues to do so throughout childhood, with some regulatory functions achieving adult-like levels by adolescence (Luna, Paulsen, Padmanabhan, & Geier, 2013; Posner, Rothbart, Sheese, & Voelker, 2014; Rothbart & Posner, 2011). While adolescents can inhibit responding as well as adults in some contexts, self-regulation within certain rewarding environments, particularly involving peers, may be more challenging and require greater effort (Anderson & Briggs, 2016; Galvan, 2013; Van Duijvenvoorde & Crone, 2013).

Individual differences in effortful and reactive self-control (impulsivity; inhibition to novelty) are evident early as 2.5 years of age with evidence for differentiation between reactive under- and over-control by age 3.5 (Eisenberg et al., 2013). Temperament traits, particularly surrounding self-control, inhibition, and disinhibition, have been related to the development of psychopathology in childhood and beyond (Kagan, 2013; Nigg, 2000, 2006).

Partly based on these developmental findings, Carver and colleagues (Carver, Johnson, & Joormann, 2008, 2014) developed a model in which the relative dominance of impulsive vs. reflective processes is modulated by serotonin, with low levels of

serotonin relating to a relative dominance of impulsive processing. An elegant aspect of this model is that it can explain effects of both trait-differences (e.g. genetic factors modulating serotonin, for example genetic variability of the serotonin transporter gene, 5HTT), and state modulations of serotonin (e.g. depletion effect through experimental manipulations or natural factors such as sleep loss). Interestingly, a recent study demonstrated that participants with a genetic variant related to relatively low levels of serotonin performed better on a sorting task under conditions that favored model-free learning, while individuals with high levels of serotonin performed better under conditions that favored model-based learning in the same task (Maddox et al., 2015).

From this perspective (Carver et al., 2008, 2014; Ernst, Pine, & Hardin, 2006), individual differences in tendencies to develop certain mental and health problems are related to the interactions of *three* broad neural systems: an impulsive approach-oriented, an impulsive avoidance-oriented, and a reflective effortful control system. This model can explain why low levels of serotonin are related both to externalizing problems (e.g., aggression) and internalizing problems (e.g., anxiety and depression). For example, strong reactive approach coupled with weak control increases chances of reactive aggression, weak approach with weak control increases chances of depressive symptoms, and strong avoidance with weak control increases chances of developing anxiety problems. Important support for this idea comes from the at first sight counter-intuitive finding that individuals who have experienced a depressive episode are also more impulsive in response to positive stimuli (Carver, Johnson, & Joormann, 2013).

As another a point in case, the famous Dunedin study, following approximately 1000 children since birth into adulthood, has demonstrated that indices of effortful control in childhood predicted later outcomes of mental and physical health (Moffitt et al., 2011), replicating and extending earlier findings by Walter Mischel and colleagues

(Mischel, Shoda, & Rodriguez, 1989). Interestingly, a recent paper based on data of the Dunedin study, argued that there could be a general factor in psychopathology (labeled the “p-factor”), in addition to the two broad classes of internalizing and externalizing behaviors (Caspi et al., 2014). One of the attractive aspects of this conceptualization is that it better fits with the developmental multiple process models described above, and indeed the p-factor was found to correlate with working memory capacity, often used as a general index of the capacity to control (Hofmann, Friese, & Wiers, 2011). In addition, the model may explain the counterintuitive positive correlation between internalizing and externalizing disorders, while in clinical practice children typically first develop either an internalizing or an externalizing problem, with occasional overlap often as a secondary diagnosis (e.g. becoming depressed as a consequence of negative social reactions to aggressive behavior). Hence, from this perspective, relatively weak effortful control and the associated tendency of reacting relatively impulsively are associated with the tendency to develop both internalizing and externalizing problems.

The next important developmental period is adolescence, the period in which most psychopathology originates (Paus, Keshavan, & Giedd, 2008), and the period in which many habits develop which impact later healthy and unhealthy life-styles (smoking, drinking, physical exercise, etc.). With the onset of puberty, testosterone rises, which has been associated with an increase in dopamine in the midbrain and at a behavioral level with sensation seeking (Steinberg, 2010) and risk taking (Peper, Koolschijn, & Crone, 2013). From the present perspective, it would suggest a shift in balance between operating systems toward more impulsive reactions especially to emotional stimuli (Carver et al., 2014), as indeed has been demonstrated both for reactions to both threat and romantic pictures (Quevedo, Benning, Gunnar, & Dahl, 2009). Adolescent risk-taking has been related to a relative immaturity of brain regions

involved in control (frontal cortices), combined with a higher reward sensitivity (Steinberg, 2010). However, changes in social and affective brain processes have also been implicated (Crone & Dahl, 2012), indicating that some of adolescents' risk-taking behaviors may be not due to a lack of ability to moderate an impulse, but rather due to other goals being more important (e.g., making a good impression on peers or potential romantic partners). Similarly, in addictive behaviors in youth, it has often been assumed that they occur because of the relative immaturity of cognitive control systems (making impulsive reactions dominant), but often also these behaviors are goal-driven (Kopetz, Lejuez, Wiers, & Kruglanski, 2013; Kopetz & Orehek, 2015). This is another example of affective goals influencing behavior through reflective rather than impulsive processes. Hence, control of impulses with an impact on addictive and other health-behaviors is not only about ability to control, but also about motivation to control (Hofmann, Friese, et al., 2008; van Deursen et al., 2015; Wiers et al., 2007).

In a series of studies on the prediction of substance use in adolescents, it was found that in adolescents with relatively weak control capacity (assessed with measures of working memory or of interference control such as the classical Stroop test), substance use and problems were better predicted by relatively automatic or implicit measures (review: Wiers, Boelema, Nikolaou, & Gladwin, 2015). These findings were observed in adolescents both in regular education (Thush et al., 2008), and in special education, both in the Netherlands (Peeters et al., 2012, 2013) and in the U.S. (Grenard et al., 2008). However, there were also some negative findings (Janssen, Larsen, Vollebergh, & Wiers, 2015), perhaps related to internet assessment. In some of the studies, the opposite pattern was found for explicit cognitive motivational measures such as alcohol-related expectancies and motives to drink, which were found to be stronger predictors in individuals with relatively strong Executive Control (EC) capacity

(Thush et al., 2008), suggesting that some adolescents (with relatively weak control capacity) are more sensitive for environmental cues to drink or smoke, while others (with stronger control capacity) make more rational decisions regarding substance use. Note that this interaction pattern was not only found for addictive behaviors, but also for other health-related behaviors, including dietary behaviors (Hofmann, Gschwendner, Friese, Wiers, & Schmitt, 2008), and aggression after alcohol (Wiers, Beckers, Houben, & Hofmann, 2009). Interestingly, from the present perspective, a similar pattern was also found in adolescents in the prediction of anxiety (Salemink & Wiers, 2012), and including the opposite pattern of stronger prediction of explicit processes in individuals with a strong control system was observed in anxiety (Salemink, Friese, Drake, Mackintosh, & Hoppitt, 2013).

Finally, it should be noted that dual process models in fact would predict a triple interaction, including not only ability to control impulses, but also motivation to do so (Hofmann, Friese, et al., 2008; Wiers et al., 2007). Indeed, in a recent study this triple interaction was partly confirmed in a sample of problem drinkers who signed up for online training: Implicit alcohol associations were especially predictive of levels of problematic drinking in those who scored low on control capacity and high on motivation to cut down (van Deursen et al., 2015). A similar pattern was also observed in a study which experimentally manipulated motivation to restrain drinking in students (who were told that greater consumption in a taste test would impair performance on a later task where they could win a reward), and found prediction of drinking by implicit associations, in those students in an ego-depletion condition (Ostafin, Marlatt, & Greenwald, 2008).¹ Hence, the overall pattern is consistent with the notion that the

¹ Note that ego-depletion has been criticized recently and appears at least to some extent to be dependent on beliefs (Job, Walton, Bernecker, & Dweck, 2013).

impact of impulsive processes on health behaviors depends on the combination of ability and motivation to control.

Implications for Interventions in Health: Training and Motivation

From the multiple processing account laid out above, health interventions can target different processes, either more impulsive or more reflective processes (Frieze, Hofmann, & Wiers, 2011), provided that there is some motivation to change the behavior. A number of training interventions have been developed which target either different relatively automatic cue-induced processes in addictive behaviors, such as an attentional bias and an approach-bias for substance-related cues, or cognitive processes related to control over impulses, such as working memory training (review: Wiers et al., 2013). The first types of training are examples of Cognitive Bias Modification (CBM), which targets domain-specific cognitive-motivational processes, such as an attentional bias, affective memory associations, or action tendencies related to disorder or health-domain specific cues. Hence, CBM in alcohol use disorders or risky drinking will contain stimuli (words or pictures) referring to alcohol, and CBM for smoking will contain cigarette-cues. In order to assess (and modify) disorder-relevant biases, it is important that the stimuli are first validated (see for a protocol: Pronk, van Deursen, Beraha, Larsen, & Wiers, 2015). The second type of training targets domain-general executive control processes, like working memory, which has indeed shown promise in relation to addictive behaviors (Bickel, Yi, Landes, Hill, & Baxter, 2011; Houben, Wiers, & Jansen, 2011), and risk for depression (Hoorelbeke, Koster, Vanderhasselt, Callewaert, & Demeyer, 2015). Another strategy concerning training of general control capacity concerns training of inhibition (Dovis, Van Der Oord, Wiers, & Prins, 2015).

One characteristic of domain-general training is that – compared to CBM – many more sessions are needed for this type of training (typically around 25, Klingberg, 2010;

while around six sessions usually suffice in CBM, Eberl et al., 2014). This makes motivation to train very important in EC-training. One way to increase this is to embed the training in a game-like program (Dovis et al., 2015). However, it should be noted in the context of substance use, that while gamification may increase motivation to do training, it does not by itself increase motivation to change the addictive behavior (Boendermaker, Prins, & Wiers, 2015). Moreover, when a boring training is advertised as a game, youth will have high expectations, based on their experience with commercial videogames, and the gamified training may briefly be less boring than the original, but subsequently be liked even less, because of a violation of expectancy effect (Boendermaker, Boffo, & Wiers, 2015).

One type of training that is sometimes misclassified from the present logic, concerns what we called selective inhibition training (Wiers et al., 2013), in which a Go/NoGo task is administered including cues of a health-related behavior (e.g. alcohol), which are systematically paired with an inhibitory (NoGo) response in the active condition, while there is no such relationship in the control condition. Because of the specific categories involved, this type of training should be regarded as a variety of CBM rather than as executive control training. Further, effects on behavior have been found after short training (a single session), and effects generalized to affective associations (Houben, Havermans, Nederkoorn, & Jansen, 2012; Houben, Nederkoorn, Wiers, & Jansen, 2011), but not to general inhibition as assessed with another standard task (Houben et al., 2012). Note that in these studies, participants (typically students) were not aware of the manipulation and did not need to have a motivation to change. Effects were short-lived (in a taste-test directly after the manipulation or in the week to follow) and weaker over time (see for a meta-analysis: Allom, Mullan, & Hagger, 2015).

However, one study successfully used this paradigm to support overweight people to

reduce weight (Veling, Koningsbruggen, Aarts, & Stroebe, 2014). In this study, volunteers in a study investigating different psychological interventions to facilitate dieting, were randomized to either active Go/NoGo training (combining food and drink pictures with a NoGo signal) or control Go/NoGo training (without food and drink pictures). In addition, participants also received an active or control implementation intention condition. In the active condition, participants were instructed to make detailed if...then plans concerning different meals, and how to handle the situation in line with the diet-goals. Hence, participants received Go/NoGo training (real/sham) and implementation intention training (real/sham). Both interventions had a positive effect on success in dietary restraint (on weight reduction), and only the effect of implementation intentions were moderated by the strength of the dietary goal, suggesting that this intervention strengthens the impact of reflective or goal-directed processes, while GoNoGo training appears to be more independent of these processes. Note that in the domain of anxiety, there is also a 'mixture' type of training. In order to effectively improve cognitive control over emotional information processing, a working memory training in the context of emotional stimuli was developed and improved cognitive control over affective information and emotion regulation stimuli, while a working memory training with neutral stimuli did not (Schweizer, Grahn, Hampshire, Mobbs, & Dalgleish, 2013; Schweizer, Hampshire, & Dalgleish, 2011). A recent study with adolescents was unable to replicate these findings, but participants were unselected adolescents who thus were not selected on motivation for change (de Voogd, Wiers, Zwitser, & Salemink, 2016).

Clinical samples are typically at least somewhat motivated to change and receive therapy in which alternative goals are activated that are incompatible with continued excessive alcohol or drug use (Wiers et al., 2016). In this context, varieties of CBM

training have been found to generate positive effects. For example, re-training of an approach-bias for alcohol was found to increase abstinence rates one year after treatment discharge by approximately 10% (Eberl et al., 2013; Wiers, Eberl, Rinck, Becker, & Lindenmeyer, 2011), with the clinical effect being mediated by the change in approach bias (Eberl et al., 2013). Re-training an attentional bias for alcohol in patients also had clinically relevant effects (Schoenmakers et al., 2010). These findings contrast with more modest or null-findings in student-populations, typically not motivated to change (Christiansen, Schoenmakers, & Field, 2015; Lindgren et al., 2015). Hence, an important issue concerning the application of CBM to the domain of health is that participants should first be at least somewhat motivated to change for CBM to exert effects. For example, a study in community smokers unmotivated to quit was successful in changing an attentional bias for cigarettes, but did not result in a change in smoking behavior (Kerst & Waters, 2014).

Increasing Motivation to Change

Motivation to change can be effectively increased through motivational interviewing (MI), a technique originally developed in addiction treatment research and now used more widely to increase motivation to change behaviors related to (mental) health (Miller & Rollnick, 2013). Tailored varieties have been developed (combined with CBT elements), to increase motivation to change and motivation to perform the training for alcohol-dependent outpatients (Boffo, Pronk, Wiers, & Mannarini, 2015), and for smoking youth (Kong et al., 2015). Another strategy can be to select people with at least some level of motivation to change. A recent study on smoking cessation provides a point in case. Potential participants signed up at an anti-smoking website for free online training to support their quit-attempt. Importantly, the quit-attempt was verified and only those people who made an actual quit attempt were selected to participate in the

training-study and randomized to real attentional re-training or a placebo variety including the same pictures. Results in this sample demonstrated significantly more success in abstinence half a year after the quit attempt after real training compared with placebo-training (Elfeddali, de Vries, Bolman, Pronk, & Wiers, 2016). These findings contrast with largely negative earlier trials in smoking (Christiansen et al., 2015), and with an online trial in problem drinkers who did not want to quit but wanted to reduce drinking, and succeeded irrespective of training condition (Wiers, Houben, et al., 2015). Before we further discuss the important question for health psychology how to influence people who are not motivated to change, it is worth pointing out that CBM also has potential in treating internalizing disorders and related health problems. Training-induced changes in attentional bias have relatively consistently been associated with changes in anxiety or emotional responsiveness to stressors (Macleod & Clarke, 2015; Van Bockstaele et al., 2014). In a similar vein, experimentally induced changes in interpretation bias (i.e., the tendency to interpret a neutral or ambiguous stimulus as being threatening) have also been shown to reduce anxiety (Hirsch, Meeten, Krahe, & Reeder, 2016). Also in this domain, there are some indications that motivation plays an important role. For example, findings regarding interpretation bias re-training appear to be more promising in a clinical sample. Note that there are also negative findings which led some to conclude that CBM is ineffective (Cristea, Kok, & Cuijpers, 2015). However, a more in-depth analysis demonstrated that those studies which did not affect the cognitive bias had no effect on emotional and clinical symptoms, but those studies that did succeed in changing the bias resulted in a medium to large effect in emotional and clinical symptoms (Macleod & Clarke, 2015). Important for the domain of health psychology is that the negative studies were often conducted over the internet.

Cognitive Bias Modification (CBM) and Emotion Regulation

Improved emotion regulation is likely to be the mechanism behind these beneficial effects of CBM in anxiety. Indeed, according to the influential emotion regulation model of Gross (Gross & Thompson, 2007), two important strategies to regulate emotions are attention deployment and reappraisal. Attention deployment refers to people's ability to selectively attend to specific positive, neutral, or negative aspects of a situation, whereas reappraisal refers to people's ability to select or change the meaning that is attached to a situation. As such, attentional bias modification procedures have been argued to improve attention deployment (MacLeod & Grafton, 2014; Todd, Cunningham, Anderson, & Thompson, 2012), and interpretation bias modification procedures are likely to improve (re)appraisal skills. In line with dual process views on mental health problems, many emotion regulation researchers have distinguished between relatively automatic, implicit emotion regulation and more effortful, explicit emotion regulation (Gyurak, Gross, & Etkin, 2011; Koole & Rothermund, 2011). CBM techniques may thus prove crucial in improving implicit emotion regulation.

Unmotivated Participants: Where to Start?

As outlined above, cognitive training paradigms have rather consistently demonstrated effectiveness in participants with some motivation to change (provided that the bias was indeed successfully changed). Through techniques like MI, motivation can be increased, but to participate in MI, there should be at least some level of motivation, to... participate in MI. What to do when this is not the case, an especially thorny problem with emerging substance use problems in youth? One obvious method is simply to pay for participation and hope that more intrinsic motivation will be activated during the following brief intervention. This may have some effect in adolescents (Kong et al., 2015), but not in students (Lindgren et al., 2015). Another strategy in adolescents and young adults is to cease the opportunity, when there is at

least for some time some motivation to change, for example after an alcohol-related injury. MI has been successfully applied in this context (Monti et al., 1999), but we are not aware of interventions targeting more automatic cognitive processes in addition in this context. Short-lived effects on cognitive processes and behavior have been reported for samples (typically students) who were not motivated to change for some of the CBM-interventions mentioned above, including selective inhibition (Allom et al., 2015), approach-bias modification (Wiers, Rinck, Kordts, Houben, & Strack, 2010), and using other techniques such as evaluative conditioning (Houben, Havermans, & Wiers, 2010), formation of implementation intentions (Sheeran et al., 2013; Veling et al., 2014), mindfulness meditation (Ostafin, Bauer, & Myxter, 2012), and physical exercise (Rensburg & Taylor, 2009). Although these paradigms may nudge people toward better health outcomes, we believe that for more enduring change, activation of goals incompatible with continuation of the unhealthy behavior is crucial (Kopetz et al., 2013; Wiers et al., 2016). Hence, from this perspective, for health interventions to be effective, a start with addressing impulsive affective processes may be helpful, and could be subjected to further investigation, but some cognitive restructuring appears to be a necessary additional ingredient for long-term change.

Substance Use as a Moderator in Health Behavior

Some final words follow about substance use and prevention from the present perspective. First, substance use can be considered as one of the possible outcomes in (dual) process models of health, but it is also an important moderator. There is ample evidence that the relative balance in decision making shifts in favor of more impulsive processes as a result of intoxication, which affects also other health behaviors such as condom use (T. K. MacDonald et al., 2000) and snacking (Hofmann & Friese, 2008). In relation to dieting, alcohol has multiple effects: it may loosen dietary control and

increase reactivity to attractive high-caloric foods, and adds many hidden calories by itself (Hofmann, Förster, Stroebe, & Wiers, 2011). This shift in control appears to be related to both strengthening of impulsive processing and weakening of EC-processes (Field, Wiers, Christiansen, Fillmore, & Verster, 2010). Further, there is some evidence for an increase in impulsive personality traits as a result of binge-drinking in adolescents (White et al., 2011), which may in the long run also affect other health behaviors, as a consequence of increased impulsivity.

Prevention

Finally, from the present perspective, traditional prevention methods may be effective, to the extent that they activate goals incompatible with initiation of the health-endangering behavior. However, the danger of non-intended effects should not be underestimated (Krank et al., 2010), and effects of general school-based prevention efforts in the substance use area are often small at best (Tobler et al., 2000). Two alternatives deserve brief mentioning. First, in (young) children an alternative could be to stimulate a relatively reflective way of processing (Diamond, 2012), and self-control training in school-children has been demonstrated to delay onset of alcohol and cigarette use years later (Lier, Huizink, & Crijnen, 2009). Second, Conrod and colleagues developed a personality-based targeted intervention approach in adolescents, where adolescents scoring high on personality traits related to increased risk for substance use and related problems (impulsivity, sensation seeking, anxiety sensitivity and hopelessness) receive a brief group-based CBT program, which teaches them different ways to deal with their needs than alcohol and drug use (e.g. participate in exciting sports in sensation seekers). Long-term effects have been found across different samples (Conrod, Castellanos-Ryan, & Strang, 2015), with specific outcomes depending on the targeted personality type (e.g. strongest effects on binge-drinking in sensation

seekers who are most prone to develop these behaviors).

Conclusion

In conclusion, although dual-process models have been criticized, especially when equated with specific brain processes, we argue that at a psychological level they can still be useful, for example to understand challenges to the development of a healthy lifestyle and for the development of interventions. Affective processes can influence impulsive decision making in health, but also reflective processes, when they concern affectively relevant goals. Cognitive training methods have shown some success in changing health-behaviors, but a critical variable for long-term success appears to be motivation to change. Ideally therefore, interventions should be developed that nudge people not motivated to change toward motivation to change and then help them (if necessary) to overcome cue-induced interfering processes with long-term healthy behaviors.

Acknowledgement

The authors wish to thank Will Cunningham, Jay van Bavel, Michael Frank, Ron Dahl and Thomas Gladwin for comments on an earlier version of the chapter.

References

- Ahadi, S. A., & Rothbart, M. K. (1994). Temperament, development, and the Big Five. *The Developing Structure of Temperament and Personality from Infancy to Adulthood*, 189-207.
- Allom, V., Mullan, B., & Hagger, M. (2015). Does inhibitory control training improve health behaviour? A meta-analysis. *Health Psychology Review*, 7199(June 2015), 1-38. <http://doi.org/10.1080/17437199.2015.1051078>

- Anderson, K. G., & Briggs, K. E. L. (n.d.). Self-regulation and decision making. In S. A. Brown & R. Zucker (Eds.), *Oxford Handbook of Adolescent Substance Abuse*. NY: Oxford University Press.
- Anderson, K. G., & Briggs, K. E. L. (2016). Self-regulation and decision making. In S. A. Brown & R. A. Zucker (Eds.), *Oxford Handbook of Adolescent Substance Abuse*. NY: Oxford University Press.
<http://doi.org/10.1093/oxfordhb/9780199735662.013.021>
- Ashby, F. G., Alfonso-Reese, L. A., Turken, U., & Waldron, E. M. (1998). A neuropsychological theory of multiple systems in category learning. *Psychological Review*, *105*(3), 442–481. <http://doi.org/10.1037/0033-295X.105.3.442>
- Bechara, A. (2005). Decision making, impulse control and loss of willpower to resist drugs: a neurocognitive perspective. *Nature: Neuroscience*, *8*(11), 1458–1463.
<http://doi.org/Doi 10.1038/Nn1584>
- Beevers, C. G. (2005). Cognitive vulnerability to depression: A dual process model. *Clinical Psychology Review*, *25*(7), 975–1002.
<http://doi.org/10.1016/j.cpr.2005.03.003>
- Bickel, W. K., Yi, R., Landes, R. D., Hill, P. F., & Baxter, C. (2011). Remember the future: Working memory training decreases delay discounting among stimulant addicts. *Biological Psychiatry*, *69*(3), 260–265.
<http://doi.org/10.1016/j.biopsych.2010.08.017>
- Boendermaker, W. J., Boffo, M., & Wiers, R. W. (2015). Exploring Elements of Fun to Motivate Youth to Do Cognitive Bias Modification. *Games for Health Journal*, *4*(6), 434–443. <http://doi.org/10.1089/g4h.2015.0053>
- Boendermaker, W. J., Prins, P. J. M., & Wiers, R. W. (2015). Cognitive Bias Modification for adolescents with substance use problems - Can serious games help? *Journal of*

Behavior Therapy and Experimental Psychiatry, 49, 13–20.

<http://doi.org/10.1016/j.jbtep.2015.03.008>

Boffo, M., Pronk, T., Wiers, R. W., & Mannarini, S. (2015). Combining cognitive bias modification training with motivational support in alcohol dependent outpatients: study protocol for a randomised controlled trial. *Trials*, 16(1), 63.

<http://doi.org/10.1186/s13063-015-0576-6>

Carver, C. S., Johnson, S. L., & Joormann, J. (2008). Serotonergic function, two-mode models of self-regulation, and vulnerability to depression: what depression has in common with impulsive aggression. *Psychological Bulletin*, 134(6), 912–43.

<http://doi.org/10.1037/a0013740>

Carver, C. S., Johnson, S. L., & Joormann, J. (2013). Major depressive disorder and impulsive reactivity to emotion: Toward a dual-process view of depression. *British Journal of Clinical Psychology*, 52(3), 285–299. <http://doi.org/10.1111/bjc.12014>

Carver, C. S., Johnson, S. L., & Joormann, J. (2014). Dual Process Models and Serotonergic Functioning. In J. P. Forgas & E. Harmon-Jones (Eds.), *Motivation and Its Regulation: The Control Within* (Vol. 16, pp. 55–78). Psychology Press.

Caspi, A., Houts, R. M., Belsky, D. W., Goldman-Mellor, S. J., Harrington, H., Israel, S., ... Moffitt, T. E. (2014). The p factor: One general psychopathology factor in the structure of psychiatric disorders? *Clinical Psychological Science*, 2(2), 119–137.

<http://doi.org/10.1177/2167702613497473>

Christiansen, P., Schoenmakers, T. M., & Field, M. (2015). Less than meets the eye: Reappraising the clinical relevance of attentional bias in addiction. *Addictive Behaviors*, 44, 43–50. <http://doi.org/10.1016/j.addbeh.2014.10.005>

Collins, A. G. E., & Frank, M. J. (2013). Cognitive control over learning: creating, clustering, and generalizing task-set structure. *Psychological Review*, 120(1), 190–

229. <http://doi.org/10.1037/a0030852>
- Conrod, P. J., Castellanos-Ryan, N., & Strang, J. (2015). Brief, Personality-Targeted Coping Skills Interventions and Survival as a Non-Drug User Over a 2-Year Period During Adolescence. *Archives of General Psychiatry*, *67*(1), 85–93.
- Cristea, I. A., Kok, R. N., & Cuijpers, P. (2015). Efficacy of cognitive bias modification interventions in anxiety and depression: Meta-analysis. *British Journal of Psychiatry*, *206*(1), 7–16. <http://doi.org/10.1192/bjp.bp.114.146761>
- Crone, E. a, & Dahl, R. E. (2012). Understanding adolescence as a period of social-affective engagement and goal flexibility. *Nature Reviews. Neuroscience*, *13*(9), 636–50. <http://doi.org/10.1038/nrn3313>
- Cunningham, W. A., & Zelazo, P. D. (2007). Attitudes and evaluations: a social cognitive neuroscience perspective. *Trends in Cognitive Sciences*, *11*(3), 97–104. <http://doi.org/10.1016/j.tics.2006.12.005>
- Daw, N. D., Niv, Y., & Dayan, P. (2005). Uncertainty-based competition between prefrontal and dorsolateral striatal systems for behavioral control. *Nature Neuroscience*, *8*(12), 1704–1711. <http://doi.org/10.1038/nn1560>
- de Voogd, L. E., Wiers, R. W., Zwitser, R. J., & Salemink, E. (2016). Emotional working memory training as an online intervention for adolescent anxiety and depression: A randomised controlled trial. *Australian Journal of Psychology*, *68*(3), 228–238.
- Deutsch, R., Gawronski, B., & Strack, F. (2006). At the boundaries of automaticity: negation as reflective operation. *Journal of Personality and Social Psychology*, *91*(3), 385–405. <http://doi.org/10.1037/0022-3514.91.3.385>
- Diamond, A. (2012). Executive Functions. <http://doi.org/10.1146/annurev-psych-113011-143750>
- Dovis, S., Van Der Oord, S., Wiers, R. W., & Prins, P. J. M. (2015). Improving executive

- functioning in children with ADHD: Training multiple executive functions within the context of a computer game. A randomized double-blind placebo controlled trial. *PLoS ONE*, *10*(4), 1–30. <http://doi.org/10.1371/journal.pone.0121651>
- Eberl, C., Wiers, R. W., Pawelczack, S., Rinck, M., Becker, E. S., & Lindenmeyer, J. (2013). Approach bias modification in alcohol dependence: do clinical effects replicate and for whom does it work best? *Developmental Cognitive Neuroscience*, *4*, 38–51. <http://doi.org/10.1016/j.dcn.2012.11.002>
- Eberl, C., Wiers, R. W., Pawelczack, S., Rinck, M., Becker, E. S., & Lindenmeyer, J. (2014). Implementation of approach bias re-training in alcoholism-how many sessions are needed? *Alcoholism, Clinical and Experimental Research*, *38*(2), 587–94. <http://doi.org/10.1111/acer.12281>
- Eisenberg, N., Edwards, A., Spinrad, T. L., Sallquist, J., Eggum, N. D., & Reiser, M. (2013). Are effortful and reactive control unique constructs in young children? *Developmental Psychology*, *49*(11), 2082.
- Eisenberg, N., Smith, C. L., & Spinrad, T. L. (2010). Effortful control: Relations with emotion regulation, adjustment, and socialization in childhood. In K. D. Vohs & R. F. Baumeister (Eds.), *Handbook of self-regulation: Research, theory, and applications*, 2nd ed. (pp. 265–282). NY: Guilford.
- Elfeddali, I., de Vries, H., Bolman, C., Pronk, T., & Wiers, R. W. (2016). A randomized controlled trial of Web-based Attentional Bias Modification to help smokers quit. *Health Psychology: Official Journal of the Division of Health Psychology, American Psychological Association*, *35*(8), 870–80. <http://doi.org/10.1037/hea0000346>
- Ernst, M., Pine, D. S., & Hardin, M. (2006). Triadic model of the neurobiology of motivated behavior in adolescence. *Psychological Medicine*, *36*(September 2005), 299–312. <http://doi.org/10.1017/S0033291705005891>

- Evans, J. S. B. T. (2008). Dual-processing accounts of reasoning, judgment, and social cognition. *Annual Review of Psychology*, *59*, 255–278.
<http://doi.org/10.1146/annurev.psych.59.103006.093629>
- Evans, J. S. B. T., & Stanovich, K. E. (2013). Dual-process theories of higher cognition: Advancing the debate. *Perspectives on Psychological Science*, *8*(3), 223–241.
<http://doi.org/10.1177/1745691612460685>
- Field, M., Wiers, R. W., Christiansen, P., Fillmore, M. T., & Verster, J. C. (2010). Acute Alcohol Effects on Inhibitory Control and Implicit Cognition : Implications for Loss of Control Over Drinking, *34*(8), 1–7. <http://doi.org/10.1111/j.1530-0277.2010.01218.x>
- Frank, M. J., Cohen, M. X., & Sanfey, A. G. (2009). Multiple systems in decision making: A neurocomputational perspective. *Current Directions in Psychological Science*, *18*(2), 73–77. <http://doi.org/10.1111/j.1467-8721.2009.01612.x>
- Friese, M., Hofmann, W., & Wiers, R. W. (2011). On taming horses and strengthening riders : Recent developments in research on interventions to improve self-control in health behaviors On taming horses and strengthening riders : Recent developments in research on interventions to improve self-control i, (937777651).
<http://doi.org/10.1080/15298868.2010.536417>
- Galvan, A. (2013). The Teenage Brain Sensitivity to Rewards. *Current Directions in Psychological Science*, *22*(2), 88–93.
- Gawronski, B., & Bodenhausen, G. V. (2006). Associative and propositional processes in evaluation: Conceptual, empirical, and metatheoretical issues: Reply to Albarracín, Hart, and McCulloch (2006), Kruglanski and Dechesne (2006), and Petty and Briñol (2006). *Psychological Bulletin*, *132*(5), 745–750. <http://doi.org/10.1037/0033-2909.132.5.745>

- Gladwin, T. E., Figner, B., Crone, E. a, & Wiers, R. W. (2011). Addiction, adolescence, and the integration of control and motivation. *Developmental Cognitive Neuroscience*, 1(4), 364–76. <http://doi.org/10.1016/j.dcn.2011.06.008>
- Gray, J. A. (1994). Framework for a taxonomy of psychiatric disorder. In S. H. M. van Goozen, N. E. Van de Poll, & J. A. Sergeant (Eds.), *Emotions: Essays on emotion theory* (pp. 29–59). Hillsdale, NJ: Lawrence Erlbaum Associates.
- Grenard, J. L., Ames, S. L., Wiers, R. W., Thush, C., Sussman, S., & Stacy, A. W. (2008). Working memory capacity moderates the predictive effects of drug-related associations on substance use. *Psychology of Addictive Behaviors : Journal of the Society of Psychologists in Addictive Behaviors*, 22(3), 426–32. <http://doi.org/10.1037/0893-164X.22.3.426>
- Gross, J. J., & Thompson, R. A. (2007). Emotion regulation: Conceptual foundations. In J. J. Gross (Ed.), *Handbook of emotion regulation* (pp. 3–26). NY: Guilford.
- Gyurak, A., Gross, J. J., & Etkin, A. (2011). Explicit and implicit emotion regulation: a dual-process framework. *Cognition and Emotion*, 25(3), 400–412.
- Hazy, T. E., Frank, M. J., & O'Reilly, R. C. (2007). Towards an executive without a homunculus: computational models of the prefrontal cortex/basal ganglia system. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, 362(1485), 1601–13. <http://doi.org/10.1098/rstb.2007.2055>
- Heatherton, T. F., & Wagner, D. D. (2011). Cognitive neuroscience of self-regulation failure. *Trends in Cognitive Sciences*, 15(3), 132–139. <http://doi.org/10.1016/j.tics.2010.12.005>
- Hirsch, C. R., Meeten, F., Krahe, C., & Reeder, C. (2016). Resolving Ambiguity in Emotional Disorders: The Nature and Role of Interpretation Biases. *Annual Review of Clinical Psychology*, 12, 281–305.

- Hofmann, W., Förster, G., Stroebe, W., & Wiers, R. W. (2011). The great disinhibitor: alcohol, food cues, and eating behavior. In V. R. Preedy, R. R. Watson, & C. R. Martin (Eds.), *Handbook of Behavior, Food and Nutrition* (pp. 2977–2991). NY: Springer.
- Hofmann, W., & Friese, M. (2008). Impulses Got the Better of Me : Alcohol Moderates the Influence of Implicit Attitudes Toward Food Cues on Eating Behavior. *Journal of Abnormal Psychology, 117*(2), 420–427. <http://doi.org/10.1037/0021-843X.117.2.420>
- Hofmann, W., Friese, M., & Strack, F. (2009). Impulse and self-control from a dual-systems perspective. *Perspectives on Psychological Science, 4*(2), 1–15. <http://doi.org/10.1111/j.1745-6924.2009.01116.x>
- Hofmann, W., Friese, M., & Wiers, R. W. (2008). Impulsive versus reflective influences on health behavior: a theoretical framework and empirical review. *Health Psychology Review, 2*(2), 111–137. <http://doi.org/10.1080/17437190802617668>
- Hofmann, W., Friese, M., & Wiers, R. W. (2011). Impulsive processes in the self-regulation of health behaviour: theoretical and methodological considerations in response to commentaries. *Health Psychology Review, 5*(2), 162–171. <http://doi.org/10.1080/17437199.2011.565593>
- Hofmann, W., Gschwendner, T., Friese, M., Wiers, R. W., & Schmitt, M. (2008). Working memory capacity and self-regulatory behavior: toward an individual differences perspective on behavior determination by automatic versus controlled processes. *Journal of Personality and Social Psychology, 95*(4), 962–77. <http://doi.org/10.1037/a0012705>
- Hoorelbeke, K., Koster, E. H. W., Vanderhasselt, M.-A., Callewaert, S., & Demeyer, I. (2015). The influence of cognitive control training on stress reactivity and rumination in response to a lab stressor and naturalistic stress. *Behaviour Research*

- and Therapy*, 69, 1–10. <http://doi.org/10.1016/j.brat.2015.03.010>
- Houben, K., Havermans, R. C., Nederkoorn, C., & Jansen, A. (2012). Beer à no-go : learning to stop responding to alcohol cues reduces alcohol intake via reduced affective associations rather than increased response inhibition. *Addiction*, 107, 1280–1287. <http://doi.org/10.1111/j.1360-0443.2012.03827.x>
- Houben, K., Havermans, R. C., & Wiers, R. W. (2010). Learning to dislike alcohol: conditioning negative implicit attitudes toward alcohol and its effect on drinking behavior. *Psychopharmacology*, 211(1), 79–86. <http://doi.org/10.1007/s00213-010-1872-1>
- Houben, K., Nederkoorn, C., Wiers, R. W., & Jansen, A. (2011). Resisting temptation: decreasing alcohol-related affect and drinking behavior by training response inhibition. *Drug and Alcohol Dependence*, 116(1–3), 132–6. <http://doi.org/10.1016/j.drugalcdep.2010.12.011>
- Houben, K., Wiers, R. W., & Jansen, A. (2011). Getting a grip on drinking behavior: training working memory to reduce alcohol abuse. *Psychological Science*, 22(7), 968–75. <http://doi.org/10.1177/0956797611412392>
- Janssen, T., Larsen, H., Vollebergh, W. A. M., & Wiers, R. W. (2015). Longitudinal relations between cognitive bias and adolescent alcohol use. *Addictive Behaviors*, 44, 51–57. <http://doi.org/10.1016/j.addbeh.2014.11.018>
- Job, V., Walton, G. M., Bernecker, K., & Dweck, C. S. (2013). Beliefs about willpower determine the impact of glucose on self-control, 110(37), 14837–14842. <http://doi.org/10.1073/pnas.1313475110>
- Kagan, J. (2013). Temperamental contributions to inhibited and uninhibited profiles. In *The Oxford Handbook of Developmental Psychology, Vol. 2*.
- Keren, G. (2013). A Tale of Two Systems: A Scientific Advance or a Theoretical Stone

- Soup? Commentary on Evans & Stanovich (2013). *Perspectives on Psychological Science*, 8(3), 257–262. <http://doi.org/10.1177/1745691613483474>
- Keren, G., & Schul, Y. (2009). Two Is Not Always Better Than One A Critical Evaluation of Two-System Theories. *Perspectives on Psychological Science*, 4(6), 533–550. <http://doi.org/10.1111/j.1745-6924.2009.01164.x>
- Kerst, W. F., & Waters, A. J. (2014). Attentional Retraining Administered in the Field Reduces Smokers' Attentional Bias and Craving. *Health Psychology*, 33(10), 1232–1240. <http://doi.org/http://dx.doi.org/10.1037/a0035708>
- Kievit, R. A., Romeijn, J.-W., Waldorp, L. J., Wicherts, J. M., Scholte, H. S., & Borsboom, D. (2011). Mind the Gap: A Psychometric Approach to the Reduction Problem. *Psychological Inquiry*, 22(2), 67–87. <http://doi.org/10.1080/1047840X.2011.550181>
- Klingberg, T. (2010). Training and plasticity of working memory. *Trends in Cognitive Sciences*, 14(7), 317–324. <http://doi.org/10.1016/j.tics.2010.05.002>
- Kong, G., Larsen, H., Cavallo, D. A., Becker, D., Cousijn, J., Salemink, E., ... Krishnan-Sarin, S. (2015). Re-training automatic action tendencies to approach cigarettes among adolescent smokers: a pilot study. *The American Journal of Drug and Alcohol Abuse*, 41(5), 425–32. <http://doi.org/10.3109/00952990.2015.1049492>
- Koole, S. L., & Rothermund, K. (2011). “I feel better but I don’t know why”: The psychology of implicit emotion regulation. *Cognition and Emotion*, 25(3), 389–399.
- Kopetz, C. E., Lejuez, C. W., Wiers, R. W., & Kruglanski, a. W. (2013). Motivation and Self-Regulation in Addiction: A Call for Convergence. *Perspectives on Psychological Science*, 8(1), 3–24. <http://doi.org/10.1177/1745691612457575>
- Kopetz, C. E., & Orehek, E. (2015). When the End Justifies the Means: Self-Defeating Behaviors as “Rational” and “Successful” Self-Regulation. *Current Directions in*

Psychological Science, 24(5), 386–391.

<http://doi.org/10.1177/0963721415589329>

Kounios, J., & Beeman, M. (2014). The Cognitive Neuroscience of Insight. *Annu. Rev. Psychol*, 65(4), 71–93. <http://doi.org/10.1146/annurev-psych-010213-115154>

Psychol, 65(4), 71–93. <http://doi.org/10.1146/annurev-psych-010213-115154>

Krank, M. D., Ames, S. L., Grenard, J. L., Schoenfeld, T., & Stacy, A. W. (2010). Paradoxical effects of alcohol information on alcohol outcome expectancies. *Alcoholism: Clinical and Experimental Research*, 34(7), 1193–1200. <http://doi.org/10.1111/j.1530-0277.2010.01196.x>

Kriete, T., Noelle, D. C., Cohen, J. D., & O'Reilly, R. C. (2013). Indirection and symbol-like processing in the prefrontal cortex and basal ganglia. *Proceedings of the National Academy of Sciences*, 110(41), 16390–16395.

<http://doi.org/10.1073/pnas.1303547110>

Lier, P. A. C. Van, Huizink, A., & Crijnen, A. (2009). Impact of a preventive intervention targeting childhood disruptive behavior problems on tobacco and alcohol initiation from age 10 to 13 years, *100*, 228–233.

<http://doi.org/10.1016/j.drugalcdep.2008.10.004>

Lindgren, K. P., Wiers, R. W., Teachman, B. A., Gasser, M. L., Westgate, E. C., Cousijn, J., ... Neighbors, C. (2015). Attempted training of alcohol approach and drinking identity associations in us undergraduate drinkers: Null results from two studies. *PLoS ONE*, 10(8), 1–21. <http://doi.org/10.1371/journal.pone.0134642>

Luna, B., Paulsen, D. J., Padmanabhan, A., & Geier, C. (2013). The teenage brain cognitive control and motivation. *Current Directions in Psychological Science*, 22(2), 94–100.

MacDonald, K. B. (2008). Effortful control, explicit processing, and the regulation of human evolved predispositions. *Psychological Review*, 115(4), 1012–1031.

<http://doi.org/10.1037/a0013327>

- MacDonald, T. K., Fong, G. T., Zanna, M. P., Martineau, A. M., Ebel, A., Ellice, J., ... Terp-, J. (2000). Alcohol Myopia and Condom Use : Can Alcohol Intoxication Be Associated With More Prudent Behavior ? *Journal of Personality and Social Psychology*, 78(4), 605–619. <http://doi.org/10.1037//0022-3514.78.4.605>
- Macleod, C., & Clarke, P. J. F. (2015). The attentional bias modification approach to anxiety intervention. *Clinical Psychological Science*, 3(1), 58–78. <http://doi.org/10.1177/2167702614560749>
- MacLeod, C., & Grafton, B. (2014). Regulation of emotion through modification of attention. In J. J. Gross (Ed.), *Handbook of emotion regulation, 2nd Ed.* (pp. 508–528). NY: Guilford\.
- Maddox, W. T., Gorlick, M. A., Koslov, S., McGeary, J. E., Knopik, V. S., & Beevers, C. G. (2015). Serotonin Transporter Genetic Variation is Differentially Associated with Reflexive- and Reflective-Optimal Learning. *Cerebral Cortex*, bhv309. <http://doi.org/10.1093/cercor/bhv309>
- Miller, W. R., & Rollnick, G. A. (2013). *Motivational Interviewing: Helping people Change*. NY: Guilford.
- Mischel, W., Shoda, Y., & Rodriguez, M. L. (1989). Delay of gratification in children. *Science*, 244(4907), 933–938.
- Moffitt, T. E., Arseneault, L., Belsky, D., Dickson, N., Hancox, R. J., Harrington, H., ... Caspi, A. (2011). A gradient of childhood self-control predicts health, wealth, and public safety. *Proceedings of the National Academy of Sciences of the United States of America*, 108(7), 2693–2698. <http://doi.org/10.1073/pnas.1010076108>
- Monti, P. M., Colby, S. M., Barnett, N. P., Spirito, A., Rohsenow, D. J., Myers, M., ... Lewander, W. (1999). Brief intervention for harm reduction with alcohol-positive older adolescents in a hospital emergency department. *Journal of Consulting and*

Clinical Psychology, 67(6), 989.

Nigg, J. T. (2000). On inhibition/disinhibition in developmental psychopathology: views from cognitive and personality psychology and a working inhibition taxonomy.

Psychological Bulletin, 126(2), 220.

Nigg, J. T. (2006). Temperament and developmental psychopathology. *Journal of Child Psychology and Psychiatry and Allied Disciplines*, 47(3-4), 395-422.

<http://doi.org/10.1111/j.1469-7610.2006.01612.x>

Ostafin, B. D., Bauer, C., & Myxter, P. (2012). Mindfulness Decouples the Relation between Automatic Alcohol Motivation and Heavy Drinking, *31(7)*, 729-745.

Ostafin, B. D., Marlatt, G. A., & Greenwald, A. G. (2008). Drinking without thinking : An implicit measure of alcohol motivation predicts failure to control alcohol use.

Behaviour Research and Therapy, 46(11), 1210-1219.

<http://doi.org/10.1016/j.brat.2008.08.003>

Ouimet, A. J., Gawronski, B., & Dozois, D. J. A. (2009). Cognitive vulnerability to anxiety: A review and an integrative model. *Clinical Psychology Review*, 29(6), 459-470.

<http://doi.org/10.1016/j.cpr.2009.05.004>

Paus, T., Keshavan, M., & Giedd, J. N. (2008). Why do many psychiatric disorders emerge during adolescence? *Nature Reviews. Neuroscience*, 9(12), 947-957.

<http://doi.org/10.1038/nrn2513>

Peeters, M., Monshouwer, K., van de Schoot, R. A. G. J., Janssen, T., Vollebergh, W. A. M., & Wiers, R. W. (2013). Automatic processes and the drinking behavior in early adolescence: A prospective study. *Alcoholism: Clinical and Experimental Research*, 37(10), 1737-1744. <http://doi.org/10.1111/acer.12156>

Peeters, M., Wiers, R. W., Monshouwer, K., van de Schoot, R., Janssen, T., & Vollebergh, W. a M. (2012). Automatic processes in at-risk adolescents: the role of alcohol-

- approach tendencies and response inhibition in drinking behavior. *Addiction (Abingdon, England)*, 107(11), 1939–46. <http://doi.org/10.1111/j.1360-0443.2012.03948.x>
- Peper, J. S., Koolschijn, P. C. M. P., & Crone, E. A. (2013). Development of risk taking: contributions from adolescent testosterone and the orbito-frontal cortex. *Journal of Cognitive Neuroscience*, 25(12), 2141–2150.
- Pessoa, L. (2008). On the relationship between emotion and cognition. *Nature Reviews Neuroscience*, 9(2), 148–58. <http://doi.org/10.1038/nrn2317>
- Posner, M. I., Rothbart, M. K., Sheese, B. E., & Voelker, P. (2014). Developing attention: Behavioral and brain mechanisms. *Advances in Neuroscience*, 2014.
- Pronk, T., van Deursen, D. S., Beraha, E. M., Larsen, H., & Wiers, R. W. (2015). Validation of the Amsterdam Beverage Picture Set: A Controlled Picture Set for Cognitive Bias Measurement and Modification Paradigms. *Alcoholism: Clinical and Experimental Research*, 39(10), 2047–2055. <http://doi.org/10.1111/acer.12853>
- Quevedo, K. M., Benning, S. D., Gunnar, M. R., & Dahl, R. E. (2009). The onset of puberty: effects on the psychophysiology of defensive and appetitive motivation. *Development and Psychopathology*, 21(1), 27–45. <http://doi.org/10.1017/S0954579409000030>
- Rensburg, K. J. Van, & Taylor, A. (2009). Acute exercise modulates cigarette cravings and brain activation in response to smoking-related images : an fMRI study. *Psychopharmacology*, 203, 589–598. <http://doi.org/10.1007/s00213-008-1405-3>
- Rothbart, M. K., & Posner, M. I. (2011). Developing mechanisms of self-regulation in early life, 3(2), 207–213. <http://doi.org/10.1177/1754073910387943>. Developing
- Salemink, E., Friese, M., Drake, E., Mackintosh, B., & Hoppitt, L. (2013). Indicators of implicit and explicit social anxiety influence threat-related interpretive bias as a

- function of working memory capacity. *Frontiers in Human Neuroscience*, 7(May), 220. <http://doi.org/10.3389/fnhum.2013.00220>
- Salemink, E., & Wiers, R. W. (2012). Adolescent threat-related interpretive bias and its modification: the moderating role of regulatory control. *Behaviour Research and Therapy*, 50(1), 40–6. <http://doi.org/10.1016/j.brat.2011.10.006>
- Satpute, A. B., & Lieberman, M. D. (2006). Integrating automatic and controlled processes into neurocognitive models of social cognition. *Brain Research*, 1079(1), 86–97. <http://doi.org/10.1016/j.brainres.2006.01.005>
- Schoenmakers, T. M., de Bruin, M., Lux, I. F. M., Goertz, A. G., Van Kerkhof, D. H. a T., & Wiers, R. W. (2010). Clinical effectiveness of attentional bias modification training in abstinent alcoholic patients. *Drug and Alcohol Dependence*, 109(1–3), 30–6. <http://doi.org/10.1016/j.drugalcdep.2009.11.022>
- Schweizer, S., Grahn, J., Hampshire, A., Mobbs, D., & Dalgleish, T. (2013). Training the emotional brain: improving affective control through emotional working memory training. *The Journal of Neuroscience*, 33(12), 5301–5311.
- Schweizer, S., Hampshire, A., & Dalgleish, T. (2011). Extending brain-training to the affective domain: Increasing cognitive and affective executive control through emotional working memory training. *PLoS ONE*, 6(9). <http://doi.org/10.1371/journal.pone.0024372>
- Sheeran, P., Gollwitzer, P. M., & Bargh, J. a. (2013). Nonconscious processes and health. *Health Psychology*, 32 VN-r(5), 460. <http://doi.org/10.1037/a0029203>
- Smith, E. R., & DeCoster, J. (2000). Dual-Process Models in Social and Cognitive Psychology: Conceptual Integration and Links to Underlying Memory Systems. *Personality and Social Psychology Review*, 4(2), 108–131. http://doi.org/10.1207/S15327957PSPR0402_01

- Steinberg, L. (2010). A dual systems model of adolescent risk-taking. *Developmental Psychobiology*, 52(3), 216–224. <http://doi.org/10.1002/dev.20445>
- Strack, F., & Deutsch, R. (2004). Reflective and impulsive determinants of social behavior. *Personality and Social Psychology Review : An Official Journal of the Society for Personality and Social Psychology, Inc*, 8(3), 220–247. http://doi.org/10.1207/s15327957pspr0803_1
- Thush, C., Wiers, R. W., Ames, S. L., Grenard, J. L., Sussman, S., & Stacy, A. W. (2008). Interactions between implicit and explicit cognition and working memory capacity in the prediction of alcohol use in at-risk adolescents. *Drug and Alcohol Dependence*, 94(1–3), 116–24. <http://doi.org/10.1016/j.drugalcdep.2007.10.019>
- Tobler, N. S., Roona, M. R., Ochshorn, P., Marshall, D. G., Streke, A. V., & Stackpole, K. M. (2000). School-based adolescent drug prevention programs: 1998 meta-analysis. *Journal of Primary Prevention*, 20(4), 275–336.
- Todd, R. M., Cunningham, W. A., Anderson, A. K., & Thompson, E. (2012). Affect-biased attention as emotion regulation. *Trends in Cognitive Sciences*, 16(7), 365–372.
- Van Bockstaele, B., Verschuere, B., Tibboel, H., De Houwer, J., Crombez, G., & Koster, E. H. W. (2014). A review of current evidence for the causal impact of attentional bias on fear and anxiety. *Psychological Bulletin*, 140(3), 682.
- van Deursen, D. S., Saleminck, E., Boendermaker, W. J., Pronk, T., Hofmann, W., & Wiers, R. W. (2015). Executive Functions and Motivation as Moderators of the Relationship Between Automatic Associations and Alcohol Use in Problem Drinkers Seeking Online Help. *Alcoholism: Clinical and Experimental Research*, 39(9), 1788–1796. <http://doi.org/10.1111/acer.12822>
- Van Duijvenvoorde, A. C. K., & Crone, E. A. (2013). The teenage brain a neuroeconomic approach to adolescent decision making. *Current Directions in Psychological Science*,

22(2), 108–113.

- Veling, H., Koningsbruggen, G. M. Van, Aarts, H., & Stroebe, W. (2014). Targeting impulsive processes of eating behavior via the internet . Effects on body weight ☆. *Appetite*, 78, 102–109. <http://doi.org/10.1016/j.appet.2014.03.014>
- White, H. R., Marmorstein, N. R., Crews, F. T., Bates, M. E., Mun, E. Y., & Loeber, R. (2011). Associations Between Heavy Drinking and Changes in Impulsive Behavior Among Adolescent Boys. *Alcoholism: Clinical and Experimental Research*, 35(2), 295–303. <http://doi.org/10.1111/j.1530-0277.2010.01345.x>
- Whiteside, S. P., & Lynam, D. R. (2003). Understanding the role of impulsivity and externalizing psychopathology in alcohol abuse: application of the UPPS impulsive behavior scale. *Experimental and Clinical Psychopharmacology*, 11(3), 210.
- Wiers, R. W., Bartholow, B. D., van den Wildenberg, E., Thush, C., Engels, R. C. M. E., Sher, K. J., ... Stacy, A. W. (2007). Automatic and controlled processes and the development of addictive behaviors in adolescents: a review and a model. *Pharmacology, Biochemistry, and Behavior*, 86(2), 263–83. <http://doi.org/10.1016/j.pbb.2006.09.021>
- Wiers, R. W., Becker, D., Holland, R., Moggi, F., Lejuez, C. W., & Yield, A. (2016). Cognitive Motivational Processes Underlying Addiction Treatment. In C. E. Kopetz & C. W. Lejuez (Eds.), *Addiction* (pp. 201–236). Psychology Press.
- Wiers, R. W., Beckers, L., Houben, K., & Hofmann, W. (2009). A short fuse after alcohol: implicit power associations predict aggressiveness after alcohol consumption in young heavy drinkers with limited executive control. *Pharmacology, Biochemistry, and Behavior*, 93(3), 300–5. <http://doi.org/10.1016/j.pbb.2009.02.003>
- Wiers, R. W., Boelema, S. R., Nikolaou, K., & Gladwin, T. E. (2015). On the Development of Implicit and Control Processes in Relation to Substance Use in Adolescence. *Current*

- Addiction Reports*, 2(2), 141–155. <http://doi.org/10.1007/s40429-015-0053-z>
- Wiers, R. W., Eberl, C., Rinck, M., Becker, E. S., & Lindenmeyer, J. (2011). Retraining automatic action tendencies changes alcoholic patients' approach bias for alcohol and improves treatment outcome. *Psychological Science*, 22(4), 490–7. <http://doi.org/10.1177/0956797611400615>
- Wiers, R. W., Gladwin, T. E., Hofmann, W., Salemink, E., & Ridderinkhof, K. R. (2013). Cognitive Bias Modification and Cognitive Control Training in Addiction and Related Psychopathology: Mechanisms, Clinical Perspectives, and Ways Forward. *Clinical Psychological Science*, 1(2), 192–212. <http://doi.org/10.1177/2167702612466547>
- Wiers, R. W., Houben, K., Fadardi, J. S., van Beek, P., Rhemtulla, M., & Cox, W. M. (2015). Alcohol Cognitive Bias Modification training for problem drinkers over the web. *Addictive Behaviors*, 40, 21–26. <http://doi.org/10.1016/j.addbeh.2014.08.010>
- Wiers, R. W., Rinck, M., Kordts, R., Houben, K., & Strack, F. (2010). Retraining automatic action-tendencies to approach alcohol in hazardous drinkers. *Addiction*, 105(2), 279–87. <http://doi.org/10.1111/j.1360-0443.2009.02775.x>
- Wiers, R. W., & Stacy, A. (2006). Implicit cognition and addiction. *Current Directions in Psychological Science*, 15, 292–296. Retrieved from <http://cdp.sagepub.com/content/15/6/292.short>